

## DOCUMENT RESUME

ED 395 963

TM 025 101

AUTHOR Holland, Paul W.  
TITLE A Note on the Covariance of the Mantel-Haenszel Log-Odds Ratio Estimator and the Sample Marginal Rates. Program Statistics Research Technical Report No. 89-85.  
INSTITUTION Educational Testing Service, Princeton, N.J.  
REPORT NO ETS-RR-89-19  
PUB DATE Feb 89  
NOTE 17p.  
PUB TYPE Reports - Evaluative/Feasibility (142)  
EDRS PRICE MF01/PC01 Plus Postage.  
DESCRIPTORS \*Difficulty Level; \*Estimation (Mathematics); \*Item Bias; \*Risk; Sampling; \*Test Items; Test Use  
IDENTIFIERS Item Bias Detection; \*Mantel Haenszel Procedure

## ABSTRACT

A simple technique, developed by A. Phillips (1987) is used to approximate the covariance between the Mantel-Haenszel log-odds-ratio estimator for a  $2 \times 2 \times k$  table and the sample marginal proportions. These results are then applied to obtain an approximate variance estimate of an adjusted risk difference based on the Mantel-Haenszel odds-ratio estimator. The adjusted risk difference is of potential value in those applications where at least one of the sample rates is descriptive of a relevant population rate. The example applies to the use of the Mantel Haenszel estimator to study the differential difficulty of test questions across groups of examinees. (Contains 11 references.) (Author/SLD)

\*\*\*\*\*  
\* Reproductions supplied by EDRS are the best that can be made \*  
\* from the original document. \*  
\*\*\*\*\*

U.S. DEPARTMENT OF EDUCATION  
Office of Educational Research and Improvement  
EDUCATIONAL RESOURCES INFORMATION  
CENTER (ERIC)

- ☒ This document has been reproduced as received from the person or organization originating it.
- ☐ Minor changes have been made to improve reproduction quality.
- Points of view or opinions stated in this document do not necessarily represent official OERI position or policy.

PERMISSION TO REPRODUCE THIS  
MATERIAL HAS BEEN GRANTED BY

H. I. BRAUN

TO THE EDUCATIONAL RESOURCES  
INFORMATION CENTER (ERIC).

RR-89-19

# A Note on the Covariance of the Mantel-Haenszel Log-Odds Ratio Estimator and the Sample Marginal Rates

Paul W. Holland



## PROGRAM STATISTICS RESEARCH

TECHNICAL REPORT NO. 89-85

BEST COPY AVAILABLE

EDUCATIONAL TESTING SERVICE  
PRINCETON, NEW JERSEY 08541

A NOTE ON THE COVARIANCE OF THE MANTEL-HAENSZEL LOG-ODDS-RATIO  
ESTIMATOR AND THE SAMPLE MARGINAL RATES

Paul W. Holland

Program Statistics Research  
Technical Report No. 89-85

Research Report No. 89-19

Educational Testing Service  
Princeton, New Jersey 08541-0001

February 1989

Copyright © 1989 by Educational Testing Service. All rights reserved.

The Program Statistics Research Technical Report Series is designed to make the working papers of the Research Statistics Group at Educational Testing Service generally available. The series consists of reports by the members of the Research Statistics Group as well as their external and visiting statistical consultants. Reproduction of any portion of a Program Statistics Research Technical Report requires the written consent of the author(s).

# ABSTRACT

A simple technique, developed in Phillips (1987), is used to approximate  $\text{Cov}(\hat{\theta}_{MH}, \hat{p}_i)$   $i = 1, 2$  where  $\hat{\theta}_{MH}$  is the Mantel-Haenszel log-odds-ratio estimator for a  $2 \times 2 \times K$  table and the  $\hat{p}_i$  are the sample marginal proportions. These results are then applied to obtain an approximate variance estimate of an adjusted risk difference based on the Mantel-Haenszel odds-ratio estimator.

## TABLE OF CONTENTS

	Page
1. Introduction.....	1
2. A Theoretical Application and Example.....	6
2.1 A general formula. ....	6
2.2 An application to a "Mantel-Haneszel adjusted risk difference".....	8
3. Discussion.....	10
References.....	11

## 1. INTRODUCTION

Consider the standard  $2 \times 2 \times K$  table whose  $k^{\text{th}}$   $2 \times 2$  layer is specified below.

	1	0	Total
Group 1	$A_k$	$B_k$	$n_{1k}$
Group 2	$C_k$	$D_k$	$n_{2k}$
Total	$m_{1k}$	$m_{0k}$	$t_k$

Probability models for the  $2 \times 2 \times K$  table include the two Binomial model (2B), Hauck, Anderson & Leahy, (1982) and the non-central (or "extended") hypergeometric model (NCH), Breslow, (1981). In both models the different  $2 \times 2$  layers of the  $2 \times 2 \times K$  table are statistically independent. In the 2B model,  $A_k$  and  $C_k$  are independent binomial variates, with  $A_k \sim B(p_{1k}, n_{1k})$  and  $C_k \sim B(p_{2k}, n_{2k})$ . In the NCH model;  $A_k$  has the non-central hypergeometric distribution given in (1.1) in which the margins  $n_{1k}$ ,  $m_{1k}$  and  $t_k$  are regarded as fixed, and  $\psi_k$  is the non-centrality parameter.

$$\text{Prob}(A_k = a | \psi_k, n_{1k}, m_{1k}, t_k) = (\psi_k)^a \binom{n_{1k}}{a} \binom{t_k - n_{1k}}{m_{1k} - a} / D, \quad (1.1)$$

where

$$D = \sum_u (\psi_k)^u \binom{n_{1k}}{u} \binom{t_k - n_{1k}}{m_{1k} - u}. \quad (1.2)$$

In (1.2), the range of summation is given by

$$\max(0, m_{1k} + n_{1k} - t_k) \leq u \leq \min(m_{1k}, n_{1k}). \quad (1.3)$$

Note that in (1.1) the integer,  $a$ , is also subject to the inequalities in (1.3).

When  $\psi_k = 1$ , (1.1) reduces to the usual hypergeometric distribution. The NCH model may be viewed as the conditional distribution of  $A_k$  in the 2B model given the total  $A_k + C_k = m_{1k}$ .

BEST COPY AVAILABLE

The common odds-ratio assumption may be expressed in both the 2B and the NCH models. In the 2B model it is expressed by assuming that the odds-ratio, i.e.,

$$\psi_k = (p_{1k}/(1-p_{1k})) / (p_{2k}/(1-p_{2k})),$$

does not depend on  $k$ , i.e.,  $\psi_k = \psi$ .

The common-odds-ratio assumption may be expressed in the NCH model by the assumption that  $\psi_k$  in (1.1) does not depend on  $k$ .

The Mantel-Haenszel (1959) estimator,  $\hat{\psi}_{MH}$ , for  $\psi$  under the common odds-ratio assumption is defined by

$$\hat{\psi}_{MH} = \left( \sum_k A_k D_k / t_k \right) / \left( \sum_k B_k C_k / t_k \right). \quad (1.4)$$

It is often convenient to work with the natural log of  $\psi$ , i.e.,

$$\theta = \ln(\psi)$$

and the corresponding estimator

$$\hat{\theta}_{MH} = \ln(\hat{\psi}_{MH}). \quad (1.5)$$

The two marginal rates (or risks) are the sample proportions:

$$\hat{p}_1 = \left( \sum_k A_k \right) / n_1,$$

and

$$\hat{p}_2 = \left( \sum_k C_k \right) / n_2, \quad (1.6)$$

where

$$n_1 = \sum_k n_{1k} \text{ and } n_2 = \sum_k n_{2k}.$$

The main results of this note are simple asymptotic expressions for the covariance between  $\hat{\theta}_{MH}$  and  $\hat{p}_1$  and  $\hat{p}_2$  that are valid for both the 2B and the NCH models. My approach is to exploit two useful formulas ((1.8) and (1.9) below)



that were developed in Phillips (1987) and given in Phillips and Holland (1987).

These are summarized in the following two lemmas.

Lemma 1: Let  $R = \sum_k A_k D_k / t_k$  and  $S = \sum_k B_k C_k / t_k$  and let  $r = E(R)$  and  $s = E(S)$  where these expectations are taken with respect to either the 2B or the NCH models. Define  $\tilde{\theta}_{MH}$  by

$$\tilde{\theta}_{MH} = \theta + \frac{R-r}{r} - \frac{S-s}{s}. \quad (1.7)$$

Then

$$\hat{\theta}_{MH} = \tilde{\theta}_{MH} + o_p\left(\frac{\text{Var}(R)}{r^2} + \frac{\text{Var}(S)}{s^2}\right). \quad (1.8)$$

The virtue of (1.8) is that it allows the non-linear  $\hat{\theta}_{MH}$  to be approximated by  $\tilde{\theta}_{MH}$  which is linear in  $R$  and  $S$ . In addition, (1.8) shows the way in which the distribution of  $\tilde{\theta}_{MH}$  approximates  $\hat{\theta}_{MH}$  --namely, that the quantity  $\frac{\text{Var}(R)}{r^2} + \frac{\text{Var}(S)}{s^2}$  must be small. Phillips and Holland (1987) show how this condition is satisfied in both the "large stratum" and "sparse data" situations.

Lemma 2: Let  $x^{(k)} = x(x-1)\dots(x-k+1)$  denote the descending factorial. If  $\alpha, \beta, \gamma, \delta$  are non-negative integers and  $\epsilon$  is any non-negative integer not exceeding  $\min(\alpha, \delta)$  then under either the 2B or the NCH model we have

$$E(A_k^{(\alpha)} B_k^{(\beta)} C_k^{(\gamma)} D_k^{(\delta)}) = \psi_k^\epsilon E(A_k^{(\alpha-\epsilon)} B_k^{(\beta+\epsilon)} C_k^{(\gamma+\epsilon)} D_k^{(\delta-\epsilon)}). \quad (1.9)$$

Equation (1.9) may be used to establish useful relationships between various covariances that involve  $A_k, B_k, C_k$  and  $D_k$ . This is illustrated in the proof of Theorem 1.

Theorem 1: Under the common odds-ratio assumption and either the 2B or the NCH model

$$(a) \text{Cov}(\tilde{\theta}_{MH}, \hat{p}_1) = 1/n_1,$$

and

$$(b) \text{Cov}(\tilde{\theta}_{MH}, \hat{p}_2) = -1/n_2.$$

From Lemma 1 we may use  $\tilde{\theta}_{MH}$  as an approximation to  $\hat{\theta}_{MH}$  and thereby use Theorem 1 to obtain asymptotic approximations to  $\text{Cov}(\hat{\theta}_{MH}, \hat{p}_1)$  and  $\text{Cov}(\hat{\theta}_{MH}, \hat{p}_2)$  under the 2B and NCH models. In section 2, I discuss an application of these covariance calculations.

Proof of Theorem 1. From the definitions made in Lemma 1,

$$\begin{aligned} \text{Cov}(\tilde{\theta}_{MH}, \hat{p}_i) &= \text{Cov}\left(\theta + \frac{R-r}{r} - \frac{S-s}{s}, \hat{p}_i\right) = \text{Cov}\left(\frac{R}{r} - \frac{S}{s}, \hat{p}_i\right) \\ &= \frac{1}{r} \left\{ \text{Cov}(R, \hat{p}_i) - \frac{r}{s} \text{Cov}(S, \hat{p}_i) \right\}. \end{aligned} \quad (1.10)$$

Hence we need expressions for  $\frac{r}{s}$ ,  $\text{Cov}(R, \hat{p}_i)$  and  $\text{Cov}(S, \hat{p}_i)$ .

It is well-known (and an easy application of Lemma 2 with  $\alpha = \delta = 1$ ,  $\beta = \gamma = 0$ , and  $\epsilon = 1$ ) that under either the 2B or the NCH models we have

$$E(A_k D_k) = \psi_k E(B_k C_k). \quad (1.11)$$

Then (1.11) may be used to show that under the common odds-ratio assumption

$$r = E(R) = \psi E(S) = \psi s,$$

or

$$\frac{r}{s} = \psi. \quad (1.12)$$

Parts (a) and (b) of Theorem 1 are proved in a similar manner so I will consider only (a).

We need expressions for  $\text{Cov}(R, \hat{p}_1)$  and  $\psi \text{Cov}(S, \hat{p}_1)$ . But

$$\begin{aligned} \text{Cov}(R, \hat{p}_1) &= \sum_k \frac{1}{t_k} \text{Cov}(A_k D_k, \hat{p}_1) \\ &= \sum_{k,j} \frac{1}{t_k} \frac{1}{n_1} \text{Cov}(A_k D_k, A_j). \end{aligned} \quad (1.13)$$

But, for  $k \neq j$ , the variates are independent so that (1.13) reduces to

$$\text{Cov}(R, \hat{p}_1) = \frac{1}{n_1} \sum_k \frac{1}{t_k} \{E(A_k^2 D_k) - E(A_k D_k) E(A_k)\}. \quad (1.14)$$

Similarly we have

$$\psi \text{Cov}(S, \hat{p}_1) = \frac{1}{n_1} \sum_k \frac{1}{t_k} \{\psi E(A_k B_k C_k) - \psi E(B_k C_k) E(A_k)\}. \quad (1.15)$$

But

$$\begin{aligned} E(A_k^2 D_k) &= E(A_k(A_k-1)D_k) + E(A_k D_k) \\ &= E(A_k^{(2)} D_k) + E(A_k D_k). \end{aligned} \quad (1.16)$$

Now apply Lemma 2 to (1.16) with  $\alpha = 2$ ,  $\beta = \gamma = 0$ ,  $\delta = 1$  and  $\epsilon = 1$  and obtain

$$E(A_k^2 D_k) = \psi E(A_k B_k C_k) + E(A_k D_k). \quad (1.17)$$

Hence, from (1.17) and (1.11), (1.14) becomes

$$\text{Cov}(R, \hat{p}_1) = \frac{1}{n_1} \sum_k \frac{1}{t_k} \{\psi E(A_k B_k C_k) - \psi E(B_k C_k) E(A_k) + E(A_k D_k)\} \quad (1.18)$$

$$= \psi \text{Cov}(S, \hat{p}_1) + \frac{1}{n_1} \sum_k \frac{1}{t_k} E(A_k D_k). \quad (1.19)$$

Thus, combining (1.19) with (1.10) and (1.12) we have

$$\text{Cov}(\tilde{\theta}_{MH}, \hat{p}_1) = \frac{1}{r} \left\{ \frac{1}{n_1} \sum_k \frac{1}{t_k} E(A_k D_k) \right\} = \frac{1}{r} \frac{1}{n_1} r = \frac{1}{n_1}. \quad \underline{\text{QED}}$$

## 2. A THEORETICAL APPLICATION AND EXAMPLE

In this section we develop the necessary formulas for applying the Taylor series- or  $\delta$ -method to obtain standard errors for functions of  $\hat{\theta}_{MH}$ ,  $\hat{p}_1$  and  $\hat{p}_2$  and then apply these results and those of section 1 to obtain an approximate standard error for a particular function of  $\hat{\theta}_{MH}$  and  $\hat{p}_2$  that arises in the use of the Mantel-Haenszel estimator to study differential difficulty of test questions across groups of examinees (Holland, 1985; Holland and Thayer, 1988).

### 2.1 A general formula

Let  $\hat{f} = f(\hat{\theta}_{MH}, \hat{p}_1, \hat{p}_2)$  be a differentiable function of  $\hat{\theta}_{MH}$ ,  $\hat{p}_1$  and  $\hat{p}_2$ . The  $\delta$ -method (Bishop, Fienberg and Holland, 1975) may be used to derive the large-sample variance of  $\hat{f}$ . It is summarized below.

Theorem 2: As the variances of  $\hat{\theta}_{MH}$ ,  $\hat{p}_1$  and  $\hat{p}_2$  go to zero, the variance of  $\hat{f}$  is approximated by:

$$\begin{aligned} \text{Var}(\hat{f}) = & \left(\frac{\partial f}{\partial \theta}\right)^2 \text{Var}(\hat{\theta}_{MH}) + \left(\frac{\partial f}{\partial p_1}\right)^2 \text{Var}(\hat{p}_1) \\ & + \left(\frac{\partial f}{\partial p_2}\right)^2 \text{Var}(\hat{p}_2) + 2 \frac{\partial f}{\partial \theta} \frac{\partial f}{\partial p_1} \text{Cov}(\hat{\theta}_{MH}, \hat{p}_1) \\ & + 2 \frac{\partial f}{\partial \theta} \frac{\partial f}{\partial p_2} \text{Cov}(\hat{\theta}_{MH}, \hat{p}_2) + 2 \frac{\partial f}{\partial p_1} \frac{\partial f}{\partial p_2} \text{Cov}(\hat{p}_1, \hat{p}_2). \end{aligned} \quad (2.1)$$

From Theorem 2 the covariances derived in section 1 may be combined with variance estimates of  $\hat{\theta}_{MH}$ ,  $\hat{p}_1$  and  $\hat{p}_2$  and the covariance of  $\hat{p}_1$  and  $\hat{p}_2$  to yield an approximate standard error for  $\hat{f}$ . Robins, Breslow, and Greenland (1986) give an estimator,  $\hat{\sigma}^2(\hat{\theta}_{MH})$ , of the variance of  $\hat{\theta}_{MH}$  that is valid in a variety of asymptotic situations. This variance estimator may be expressed in the notation of Lemma 1 as

$$\hat{\sigma}^2(\hat{\theta}_{MH}) = \frac{2}{R} \sum_k t_k^{-2} [A_k D_k + \hat{\psi}_{MH} B_k C_k] [A_k + D_k + \hat{\psi}_{MH} (B_k + C_k)]. \quad (2.2)$$

This variance estimator is also discussed in Phillips and Holland (1987).

The following lemma summarized the variances and covariance of  $\hat{p}_1$  and  $\hat{p}_2$  in the 2B and NCH cases.

Lemma 3: (a) In the 2B case:

$$\text{Var}(\hat{p}_i) = \frac{1}{n_i} \sum_k w_{ik} p_{ik} (1-p_{ik}) \quad (2.3)$$

where  $w_{ik} = n_{ik}/n_i$ , and

$$\text{Cov}(\hat{p}_1, \hat{p}_2) = 0. \quad (2.4)$$

(b) In the NCH case:

$$\text{Var}(\hat{p}_i) = \frac{1}{n_i} \sum_k \text{Var}(A_k) \quad (2.5)$$

and

$$\text{Cov}(\hat{p}_1, \hat{p}_2) = - \frac{1}{n_1 n_2} \sum_k \text{Var}(A_k). \quad (2.6)$$

In Theorem 3, part (a), it is clear that estimates of the variances and covariances of  $\hat{p}_1$  and  $\hat{p}_2$  under the 2B model are straightforward. For example,  $A_k/n_{1k}$  and  $C_k/n_{2k}$  can be used as estimates of  $p_{1k}$  and  $p_{2k}$ , respectively, in (2.3). On the other hand, by Jensen's inequality we have

$$\sum_k w_{ik} p_{ik} (1-p_{ik}) \leq \bar{p}_i (1-\bar{p}_i), \quad (2.7)$$

where  $\bar{p}_i = \sum_k w_{ik} p_{ik}$ . Hence from (2.3) and (2.7) we see that the simple "binomial variance" estimate,

$$\frac{1}{n_i} \hat{p}_i (1 - \hat{p}_i), \quad (2.8)$$

provides an estimate of  $\text{Var}(\hat{p}_i)$  that is, at worst, an over-estimate. When  $K$  is large and some of the  $t_k$  are small (2.8) is often a better estimate of the variance of  $\text{Var}(\hat{p}_i)$  than the one obtained by substituting the sample proportions,  $\hat{p}_{ik}$ , for  $p_{ik}$  in (2.3).

From part (b) of Lemma 3 it is evident that estimates of the variances and covariances of  $\hat{p}_1$  and  $\hat{p}_2$  under the NCH model all require estimates of  $\sum_k \text{Var}(A_k)$ , which, in turn, involves the estimates of the variance of the NCH variate,  $A_k$ . Harkness (1965) discusses the moments of  $A_k$  in the NCH case. I do not know whether or not those results can be used to give valid estimates of  $\sum_k \text{Var}(A_k)$  that are needed to use the NCH part of Lemma 3.

## 2.2 An application to a "Mantel-Haenszel adjusted risk difference".

In biomedical applications,  $\hat{\psi}_{MH}$  provides an adjusted estimate of the relative odds of getting a disease in an exposed group of individuals compared to an unexposed group. The adjustment is for differences in the distribution of potential confounding variables that may exist between the two groups.

Holland (1985) discussed the use of  $\hat{\theta}_{MH}$  as an adjusted measure of "bias" in test questions. In this use of the  $\hat{\theta}_{MH}$ , "getting the disease" is replaced by "getting the test item right" and the "exposed" and "unexposed" groups are replaced by a reference and a focal group of examinees (i.e. White and Blacks or Males and Females). The adjustment is for overall test performance. Since that suggestion, the use of the Mantel-Haenszel procedure to measure "item bias" has become wide-spread at testing organizations such as Educational Testing Service. In these testing applications, there is an interest in expressing the estimated logit differences,  $\hat{\theta}_{MH}$ , in terms of the probability scale as an adjusted difference in proportions (eg. Dorans and Kulick, 1986). One way of expressing  $\hat{\theta}_{MH}$  in the "p-scale" is the following statistic that in biometric terms might be called the "Mantel-Haenszel adjusted risk difference",

$$RD_{MH} = \hat{p}_2 - \frac{\hat{p}_2 \exp\{\hat{\theta}_{MH}\}}{(1-\hat{p}_2) + \hat{p}_2 \exp\{\hat{\theta}_{MH}\}} \quad (2.9)$$

The second term in the right-hand-side of (2.9) is the value of  $p_1$  that would be obtained if  $\hat{p}_2$  were "adjusted" by  $\hat{\theta}_{MH}$ , i.e. if  $p_1$  were chosen to solve for  $p_1$  in the equation,

$$\hat{\theta}_{MH} = \log\left(\frac{p_1}{1-p_1} / \frac{\hat{p}_2}{1-\hat{p}_2}\right). \quad (2.10)$$

The raw risk difference,  $\hat{p}_2 - \hat{p}_1$ , makes no use of the matching or stratification that is available and, for this reason, is not of much practical value except in special circumstances. The Mantel Haenszel adjusted risk difference is based on the stratification and is therefore a type of "standardized" risk difference.

If  $RD_{MH}$  from (2.9) is used, the need for its standard error arises and the results of sections 1 and 2.1 may be used to obtain an estimate of the variance of  $RD_{MH}$  under the 2B model. This is given in Theorem 3, below.

Theorem 3: Under the 2B model, the common odds-ratio assumption and any conditions that insure the approximation of  $\hat{\theta}_{MH}$  by  $\tilde{\theta}_{MH}$ , the variance of  $RD_{MH}$  in (2.9) is estimated by

$$\frac{1}{n_2} (1-G)^2 \hat{p}_2 (1-\hat{p}_2) + G^2 [\hat{p}_2(1-\hat{p}_2)]^2 \hat{\sigma}^2(\hat{\theta}_{MH}) + \frac{2}{n_2} G(1-G) \hat{p}_2(1-\hat{p}_2) \quad (2.13)$$

where

$$G = \hat{\psi}_{MH} / (1-\hat{p}_2 + \hat{\psi}_{MH} \hat{p}_2)^2$$

and  $\hat{\sigma}^2(\hat{\theta}_{MH})$  is the Robins-Breslow-Greenland variance estimate of  $\hat{\theta}_{MH}$  given in (2.2).

Proof: Let  $f(\hat{\theta}_{MH}, \hat{p}_1, \hat{p}_2)$  be defined by  $RD_{MH}$  in (2.9). Then the relevant derivatives from Theorem 2 are easily shown to be

$$\frac{\partial f}{\partial \theta} = -p_2(1-p_2)G, \quad \frac{\partial f}{\partial p_1} = 0, \quad \text{and} \quad \frac{\partial f}{\partial p_2} = 1 - G. \quad (2.14)$$

These derivatives when combined, via Theorem 2, with the covariance in Theorem 1b, and using (2.8) instead of (2.3) yields the result. QED.

### 3. DISCUSSION

The technique used to prove Theorem 1 (i.e. Lemma 1 and 2) is useful in its own right since it is simple and yet widely applicable to computations involving the Mantel-Haenszel estimator. In addition, I think it is rather remarkable that the asymptotic covariances of  $\log(\hat{\Psi}_{MH})$  and the  $\hat{p}_i$  are as simple as they appear in Theorem 1.

The adjusted risk difference,  $RD_{MH}$ , is of potential value in those applications where at least one of the sample rates, say  $\hat{p}_2$ , is descriptive of a relevant population rate. This occurs in the testing applications referred to earlier but may also arise in prospective epidemiological studies as well. The variance estimate in Theorem 3 is asymptotically valid whenever the Robins-Breslow-Greenland estimate of the variance of  $\hat{\theta}_{MH}$  is valid with the added proviso that the "binomial variance estimate", (2.8), be an appropriate estimate of the variance of  $\hat{p}_2$ . Thus the variance estimate in Theorem 3 will be most useful in the so-called "sparse-data" case where  $K$  is large and the  $t_k$  are not. In the large stratum case, i.e. when  $K$  is small and the  $t_k$  are large, it may be better to use formula (2.3) to estimate the variance of  $\hat{p}_2$ . This substitution would only change the first term of formula (2.13).

I have not performed a small sample study of the behavior of the variance estimate in Theorem 3, but because of its close connection to the Robins-Breslow-Greenland variance estimate for  $\hat{\theta}_{MH}$  I would expect it to perform quite well in both the sparse data and the large stratum cases.

To extend Theorem 3 to the NCH model it would be necessary to have a useful estimate of  $\sum_k \text{Var}(A_k)$  for the NCH case in order to apply Lemma 2(b). I do not know of any results in this area. However, in the NCH model, while the sample marginal rates  $\hat{p}_1$  and  $\hat{p}_2$  are still defined by (1.6), it is not clear what meaning to attach to them as estimates of population rates. Thus,  $RD_{MH}$  may not be a useful parameter in the NCH case.



## REFERENCES

- Bishop, Y. M. M., Fienberg, S. & Holland, P. W. (1975) Discrete multivariate analysis: theory and practice. Cambridge, MA: M.I.T. Press.
- Breslow, N. (1981) Odds-ratio estimators when the data are sparse. Biometrika, 68, 73-84.
- Dorans, N. J. & Kulick, E. (1986) Demonstrating the utility of the standardization approach to assessing unexpected differential item performance on the Scholastic Aptitude Test. Journal of Educational Measurement, 23, 355-368.
- Harkness, W. L. (1965) Properties of the extended hypergeometric distribution. Annals of Mathematical Statistics, 36, 938-945.
- Hauck, W. W., Anderson, S. & Leahy III, F. J. (1982) Finite-sample properties of some old and some new estimators of a common odds ratio from multiple 2x2 tables. Journal of the American Statistical Association, 77, 145-152.
- Holland, P. W. (1985) On the study of differential item performance without IRT. Proceedings of the Military Testing Association, October.
- Holland, P. W. & Thayer, D. T. (1988) Differential item performance and the Mantel-Haenszel procedure. In Wainer, H. and Braun, H. (Eds.) Test Validity. Hillsdale, NJ: Lawrence Erlbaum Associates.
- Mantel, N. & Haenszel W. (1959) Statistical aspects of the analysis of data from retrospective studies of disease. Journal of the National Cancer Institute, 22, 719-748.
- Phillips, A. (1987) A Study of the variance estimators of the Mantel-Haenszel log-odds-ratio estimate. Ph.D. dissertation, Department of Mathematics and Statistics, University of Windsor, Windsor, Ontario.
- Phillips, A. & Holland, P. W. (1987) Estimation of the variance of the Mantel-Haenszel log-odds-ratio estimate. Biometrics, 43, 425-431.
- Robins, J., Breslow, N. & Greenland, S. (1986) Estimators of the Mantel-Haenszel variance consistent in both sparse data and large-strata limiting models. Biometrics, 42, 311-323.